

**PERFORMANCE DATA SHEET**

2131

**Monoclonal anti-human CD80 (B7-1)\* (IgG1 isotype)****mAb name/Clone:** P1.H5.A1.A1**Isotype:** Mouse IgG1 $\kappa$ **Immunogen:** Human CD80 (B7-1) Ig Fusion Protein**CATALOG#:** 110-820 (Preservative free)**QUANTITY:** 100  $\mu$ g**CONCENTRATION:** 1.0 mg/ml

**INFORMATION:** Human CD80 (B7-1) is a costimulatory ligand for CD28 and CTLA-4. CD80 is expressed on activated B cells. Antibody P1.H5.A1.A1 recognizes the CD80 molecule and blocks binding of soluble CD152 Ig fusion protein to CD80.

**References:** C.B. Thompson, (1995) Cell **81**: 979-982. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 682-684.

**STORAGE CONDITIONS:** Store at 2 - 5°C. Open under aseptic conditions. Freeze/Thawing is not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: \_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl. Product was 0.2 $\mu$  filtered and vialled under aseptic conditions.

**PRODUCTION:** Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

**PERFORMANCE:** Five x 10<sup>5</sup> cultured human **Raji** cells were washed and incubated 45 minutes on ice with 80  $\mu$ l of anti-CD80 at 10  $\mu$ g/ml. Cells were washed twice and incubated with 2<sup>o</sup> reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.94** log<sub>10</sub> fluorescent units when compared to a Mouse IgG1 negative control (Catalog # 278-010).

*\*This Product is intended for Laboratory Research use only.*

**Binding of anti-CD80 mAb +GAM/FITC to human Raji cells**

