

PERFORMANCE DATA SHEET

3101

Monoclonal anti-human IgG/R-PE*

mAb name/Clone: ICO-97

Isotype: Mouse IgG1

Immunogen: Human Ig

CATALOG#: 139-050

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2 ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5 x 10⁵-cell /test)

INFORMATION: Human immunoglobulins are glycoproteins composed of two disulfide-bonded heavy (H) chain subunits, each of which is linked by interchain disulfide bonds to a light (L) chain forming a tetramolecular complex. There are five classes of immunoglobulins, designated IgG, IgA, IgM, IgD and IgE, which are defined by differences in the constant region of H chains. L chains are divided into kappa or lambda classifications based on structural antigenic differences. All classes of immunoglobulins have been found on the cell surface of B lymphocytes where they function as antigen receptors to elicit antigen-dependent proliferation and secretion of antigen specific soluble circulating antibodies. Antibody ICO-97 recognizes cell surface human IgG on B cells.

References: 1. Basic and Clinical Immunology, Seventh edition (D. P. Sites & A. I. Terr, eds.) Appleton & Lange., Norwalk, CT (1991).

STORAGE CONDITIONS: Store at 2 - 5°C. Do not freeze! Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at 0.5 µg/ml with a PE:Ab ratio of 0.7

PERFORMANCE: Five x 10⁵ human IgG-coated microparticles (*Inquire*) were incubated 45 minutes on ice with 80 µl of product at a dilution factor of 1:50 (10 µg/ml). They were washed three times, fixed and analyzed by FACS. Microparticles stained positive with a mean shift of 1.25 log₁₀ fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050). Binding was blocked when particles were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-huIgG antibody (Catalog #139-020).

**Research use only. Not for use in Diagnostic procedures.*

Binding of anti-human IgG/PE to human IgG1k-coated microparticles

