

PERFORMANCE DATA SHEET

1740

Monoclonal anti-human CD4*

mAb name/Clone: QS4120

Isotype: Mouse IgG1

Immunogen: CD4+ transfectant/ CEM

CATALOG#: 147-820 (Preservative Free)

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD4 is a cell surface glycoprotein of 55 kd expressed on most thymocytes, on about two thirds of peripheral T cells, and on some monocyte macrophage lineage cells. CD4 interacts with MHC Class II molecules in an accessory role during foreign antigen recognition by T cells. The extracellular portion of CD4 is an array of four Ig-like domains. The amino terminal region of CD4 is a receptor for the HIV-1 viral protein gp120. Antibody QS4120 recognizes the domain 1 region of the CD4 molecule. Antibody QS4120 will block binding of HIV-1 gp120 protein to CD4 and also blocks HLA Class II Rosette formation.

References: D. Healey, et al. (1990) J Exp Med **172**: 1233-1242. Leukocyte Typing IV (W. Knapp, et al, eds.) Oxford University Press, Oxford, (1989) p. 314-330. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 476-478. R. König, et al, (1995) J Exp Med **182**: 779-787.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin. Product was 0.2µ sterile filtered and vialled under aseptic conditions.

PERFORMANCE: Five x 10⁵ cultured **HPB-MLT** human tumor cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD4 antibody at 5 µg/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.84 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog # 278-010) at a similar concentration.

Binding of anti-CD4 antibody to human HPB-MLT tumor cells + GAM/FITC



* **Research Use Only. Not for use in Diagnostic procedures.**