

PERFORMANCE DATA SHEET

1816

Monoclonal anti-human CD10(CALLA)/FITC*

mAb Name/Clone: SN5c/L4-1A1

Isotype: Mouse IgG1 κ

Immunogen: Membrane preparation of human B-lineage leukemia cells

CATALOG#: 157-040

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2ml

WORKING DILUTION: 1:50 (or use 1.6 μ l of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Human CD10 is a well known marker for common acute lymphoblastic leukemia antigen (CALLA). CD10 is expressed on normal early B cells in bone marrow and is also on fetal thymocytes. CD10 expression is lost as the T and B lineage cells mature and only appears again on activated germinal center B cells. CD10 is present on neutrophils. CD10 is a neutral endopeptidase and hydrolyzes peptide bonds on the amino side of hydrophobic amino acids. Antibody SN5c recognizes the 100 kd membrane-associated peptidase CD10.

References: H. Matsuzaki, et al, (1987) Cancer Research **47**: 2160-2166. W.C. Biddle, et al, (1989) Leukemia Research **13**: 699-707. H. Hara, et al, (1988) Cancer Research **48**: 4673-4680. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 505-507.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____


BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at 0.25 mg/ml with a Fluorescein/IgG molar ratio of 8.9.

PERFORMANCE: Five x 10⁵ cultured Raji cells were incubated 45 minutes on ice with 80 μ l of anti-CD10/FITC at a 1:50 dilution factor (5 μ g/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.0 log₁₀ fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 μ l of 0.5 mg/ml anti-CD10 antibody (Catalog #157-020).

**This Product is intended for Laboratory Research use only.*

Binding of anti-CD10/FITC to human cell lines



Ancell Corporation P.O. Box 87 243 Third Street North Bayport, MN 55003-0087 USA
Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940