PERFORMANCE DATA SHEET

2132

Ancell

Monoclonal anti-human CD17(lactosylceramide)*

mAb name/Clone: Huly-m13/H018.3G-6.F5

Isotype: Mouse IgMk

Immunogen: Human beta-2 microglobulin associated proteins from a detergent lysate of PBL

CATALOG#: 166-020 QUANTITY: 100 μg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Antibodies to human CD17 react with lactosylceramide which is expressed on granulocytes, monocytes, platelets, and basophils (1). CD17 expression is downmodulated on activated granulocytes. Antibody Huly-m13 recognizes CD17 on myeloid cells.

References: (1)Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995) p. 822-823.

STORAGE CONDITIONS: *Store at 2 - 5^oC*. Do not Freeze!

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____


BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was purified to 95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin) using size exclusion chromatography.

PERFORMANCE: Five x 10^5 cultured **Nalm-6** human tumor cells were washed and preincubated 5 minutes with 20 μ l of 250 μ g/ml human IgG (to block non specific binding) after which they were incubated 45 minutes on ice

with 80 μ l of anti-CD17 antibody at 10 μ g/ml. Cells were washed twice and incubated with 2° reagent Goat anti-Mouse IgG/FITC (Catalog #232-011); this reagent cross reacts well with Mouse IgM, after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.1 \log_{10} fluorescent units when compared to a Mouse IgM negative control (Catalog #290-010).

Binding of anti-CD17 mAb +GAM/FITC to human Naim-6 cells



* Research Use Only. Not for use in Diagnostic procedures.

Ancell Corporation P.O. Box 87 Bayport, MN 55003-0087 USA Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940