

PERFORMANCE DATA SHEET

1816

Monoclonal anti-human CD18 ($\beta 2$ Integrin)/Biotin*

mAb name/Clone: IB4

Isotype: Mouse IgG2a

Immunogen: Human polymorphonuclear leukocytes

CATALOG#: 167-030

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD18 ($\beta 2$ integrin) forms a heterodimer with either α^L , α^M , or α^X integrins (CD11a, b or c). CD11/CD18 heterodimeric molecules are involved with cell/cell and cell/extracellular adhesion in immune and inflammatory responses. Antibody IB4 recognizes the $\beta 2$ integrin subunit (CD18) adhesion molecule of about 95 kd. Antibody IB4 blocks binding of ICAM-1 and ICAM-3 to LFA-1.

References: S.D. Wright et al, (1983) Proc Natl Acad Sci USA **80**: 5699-5703. R.C. Landis et al, (1994) J Cell Biol **126**: 529-537. M. Kaneko, et al, (1995) J Immunol **155**: 2631-2641. A. Mazzone & G. Ricevuti, (1995) Haematologica **80**: 161-175.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/thawing not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

PERFORMANCE: Briefly, 5×10^5 cultured HPB-MLT cells were washed and incubated 45 minutes on ice with 80 μ l of product at 5 μ g/ml. Cells were washed twice and incubated with 2^o reagent Streptavidin/R-Phycoerythrin (Catalog #253-050), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.67 log₁₀ fluorescent units when compared to a Mouse IgG2a/Biotin negative control (Catalog #281-030) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 μ l of 0.5 mg/ml anti-CD18 antibody (Catalog #167-020).

**This Product is intended for Laboratory Research use only.*

Binding of anti-CD18/Biotin to human cell lines

