PERFORMANCE DATA SHEET

2519

Monoclonal anti-human CD21 (CR2)*



mAb name/Clone: BU33 *Isotype:* Mouse IgG1

Immunogen: Human HFB-1 cell line

CATALOG#: 170-020 QUANTITY: 100 μg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD21 is a receptor for complement fragments C3d, C3dg or ic3D and also EBV. CD21 is also a ligand for CD23 and plays a role in IgE synthesis. Antibody BU33 recognizes the CD21 molecule of about 145 kd. Antibody BU33 inhibits binding to CD23.

References: Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 516-523. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 535-538. J.Y. Bonnefoy, et al, (1995) Intl Archiv Allergy Immunol 107: 40-42. A.D. Beaulieu, et al, (1995) Blood 86: 2789-2798. H. Molina, et al, (1995) J Immunol 154: 5426-5435.

STORAGE CONDITIONS: *Store at 2 - 5^oC*. Freeze/Thawing is not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10^5 cultured **HPB-MLT** cells were washed and incubated 45 minutes on ice with 80 μ l of anti-CD21 antibody at **20** μ g/ml. Cells were washed twice and incubated with 2^0 reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.47** \log_{10} fluorescent units when compared to a Mouse IgG1 negative control (Catalog # 278-010) at a similar concentration.

*Research use only. Not for use in Diagnostic procedures.

Binding of anti-CD21 mAb +GAM/FITC to human HPB-MLT cells

