

PERFORMANCE DATA SHEET

1816

Monoclonal anti-human CD27/FITC*

mAb name/Clone: M-T271

Isotype: Mouse IgG1

Immunogen: T cell line (from patient with CLL)

CATALOG#: 176-040

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5×10^5 -cell test)

INFORMATION: Human CD27 is a lymphocyte specific member of the tumor necrosis factor (TNF) receptor family and is found primarily on peripheral blood T cells and on a subpopulation of B cells and NK cells. The ligand for CD27 is CD70, which is a member of the TNF ligand superfamily. Antibody M-T271 recognizes the dimeric CD27 molecule of approximately 110 kd.

References: Leukocyte Typing IV (W. Knapp, et al, eds.) Oxford University Press, Oxford, (1989) p. 350-352.

K. Agematsu, et al, (1994) J Immunol **153**: 1421-1429. R.Q. Hintzen, et al, (1994) Immunol Today **15**: 307-311.

Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 356-360, 435-437.

K. Agematsu, et al, (1995) J Immunol **154**: 3627-3635.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN_3 (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC.

Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at 0.5 mg/ml with a Fluorescein/IgG molar ratio of 10.8.

PERFORMANCE: Five $\times 10^5$ ficoll prepared human peripheral blood lymphocytes per tube were washed and pre incubated 5 minutes with 20 µl of 250 µg/ml human IgG (to block non specific binding) after which they were washed and incubated 45 minutes on ice with 80 µl of anti-CD27/FITC at a 1:50 dilution factor (10 µg/ml). Cells were washed three times, fixed and analyzed by FACS. A net 56% subpopulation of the cells stained positive with a mean shift of 1.24 \log_{10} fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040). Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD27 antibody (Catalog #176-020).

**Research Use Only. Not for use in Diagnostic procedures.*

Binding of anti-CD27/FITC to human PBMC

