

**PERFORMANCE DATA SHEET**

2638

**Monoclonal anti- human CD31(PECAM-1)/R-PE \***

**mAb name/Clone:** 158-2B3

**Isotype:** Mouse IgG1

**Immunogen:** Stimulated human leukocytes

**CATALOG#:** 180-050

**QUANTITY:** 120 tests

**VOLUME IN VIAL:** 0.2 ml

**WORKING DILUTION:** 1:50 (or use 1.6µl of concentrated stock per 5 x 10<sup>5</sup>-cell test)

**INFORMATION:** Human CD31 is an adhesion molecule expressed on platelets, endothelial cells, leukocytes and their bone marrow precursors. CD31 plays a role in homophilic adhesion and heterophilic transendothelial migration (3). Antibody 158-2B3 reacts with domain 1 of CD31 and blocks homophilic interaction and heterophilic transendothelial migration (1).

**References:** 1) Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 362-372. 2) P.J. Newman, (1997) J Clin Invest **99**: 3-10. 3) K.L. Yong, et al, (1998) Blood **91**: 1196-1205.

**STORAGE CONDITIONS:** Store at 2 - 5°C. Do not freeze! Protect from light.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: \_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at **0.5 mg/ml** with an A<sub>565</sub>/A<sub>280</sub> ratio of 3.2.

**PERFORMANCE:** Five x 10<sup>5</sup> cultured **Jurkat** cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD31/R-PE at a **1:50** dilution (10 µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.59 log<sub>10</sub>** fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD31 antibody (Catalog #180-020).

\* **Research Use Only. Not for use in Diagnostic procedures..**

**Binding of anti-CD31/PE to human Jurkat cells**

