## PERFORMANCE DATA SHEET

1817

## Monoclonal anti-human CD36\*



*mAb name/Clone:* SMO *Isotype:* Mouse IgM

Immunogen: Human tonsil cells/PBM

**CATALOG#: 185-020 QUANTITY: 100 μg** 

**CONCENTRATION: 1.0 mg/ml** 

**INFORMATION:** Human CD36 is a major platelet glycoprotein also found on monocytes, macrophages and some endothelial cells. CD36 may play a role in cell adhesion as a receptor for thrombospondin. CD36 is a receptor for adhesion of malaria infected erythrocytes. CD36 is also proported to have a role in phagocytosis. Antibody SMO recognizes the CD36 molecule of 88 kd.

**References:** N. Hogg, et al, (1984) Immunology **53**: 753-767. P. Oquendo, et al, (1989) Cell **58**: 95-101. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1269-1289. Y. Ren, et al, (1995) J Exp Med **181**: 1857-1862.

STORAGE CONDITIONS: Store at 2 - 5°C. Do not Freeze!


**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

**PRODUCTION:** Antibody from (low FBS containing) tissue culture supernatant was purified to 95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin) using size exclusion chromatography.

**PERFORMANCE:** Five x  $10^5$  ficoll prepared **human peripheral blood mononuclear cells** were washed and pre incubated 5 minutes with 20  $\mu$ l of 250  $\mu$ g/ml human IgG (to block non specific binding) after which they were incubated 45 minutes on ice with 80 $\mu$ l of anti-CD36 antibody at a concentration of **5**  $\mu$ g/ml. Cells were washed twice and incubated with 20 reagent Goat anti-Mouse Ig/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. A net **16%** sub population of the cells stained positive with a mean shift of **1.05**  $\log_{10}$  fluorescent units when compared to a Mouse IgM negative control (Catalog #290-010).

Binding of anti-CD36 antibody to human cell types



\* Research Use Only. Not for use in Diagnostic procedures.

.