



PERFORMANCE DATA SHEET

2304

Monoclonal anti-human CD43 (leukosialin)/FITC*

mAb name/Clone: DFT1

Isotype: Mouse IgG1

Immunogen: Human KG-1 tumor cells

CATALOG#: 192-040

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5×10^5 -cell test)

INFORMATION: Human CD43 is a heavily O-glycosylated and sialylated transmembrane protein that is mainly found on T cells, but not on B cells. Antibody DFT1 recognizes epitope A of the CD43 molecule on pMN of about 135 kd (1,3). Antibody DFT1 partially induces apoptosis in hemopoetic progenitor cells and also induces hemopoetic aggregation.

Antibody DFT1 partially induces apoptosis in hemopoetic progenitor cells and also induces hemopoetic aggregation.

References: 1.) W.P. Stross, et al, (1989) J Clin Pathol 42:953-961. 2.) Leukocyte Typing IV (W. Knapp, et al, eds.) Oxford University Press, Oxford, (1989) p. 1697-1713. 3.) Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 494-497.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at 0.5 mg/ml.

PERFORMANCE: Five $\times 10^5$ cultured **Jurkat** cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD43/FITC at a 1:50 dilution factor (10µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.52 log₁₀ fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD43 antibody (Catalog #192-020).

Binding of anti-CD43/FITC to human cell lines



* **Research Use Only. Not for use in Diagnostic procedures.**

Ansell Corporation P.O. Box 87 Bayport, MN 55003-0087 USA
Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940