## PERFORMANCE DATA SHEET

1817

## Monoclonal anti-human CD45\*

*mAb name/Clone:* C11 *Isotype:* Mouse IgG2ak

Immunogen: Peripheral blood mononuclear cells

**CATALOG#: 196-020 QUANTITY: 100 μg** 

**CONCENTRATION: 1.0 mg/ml** 

**INFORMATION:** Human CD45 is expressed on all cells of hematopoietic origin, except erythrocytes. CD45 is a transmembrane tyrosine phosphatase which can exist in at least nine different isoforms resulting from tissue-specific alternative RNA splicing of exons 4-7 of a single gene coding for the various N-terminal peptide segments. CD45 plays an essential role in lymphocyte activation. Antibody C11 recognizes all isoforms of CD45. *References:* U. Dianzani, et al, (1992) Eur J Immunol **22**: 365-371. I.S. Trowbridge & M.L. Thomas, (1994) Annu Rev Immunol **12**: 85-116. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 386-399. D. Leitenberg, et al, (1996) J Exp Med **183**: 249-259.

**STORAGE CONDITIONS:** *Store at 2 - 5<sup>o</sup>C*. Freeze/Thawing is not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5mg/ml Gentamicin sulfate (as a preservative).

**PRODUCTION:** Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin.

**PERFORMANCE:** Five x 10<sup>5</sup> ficoll prepared **human peripheral blood mononuclear cells** were washed and pre incubated 5 minutes with 20 μl of 250 μg/ml human IgG (to block non specific binding) after which they were incubated 45 minutes on ice with 80 μl of anti-CD45 antibody at a concentration of **5 μg/ml**. Cells were washed twice and incubated with 2<sup>o</sup> reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **2.32** log<sub>10</sub> fluorescent units when compared to a Mouse IgG2a negative control (Catalog #281-010) at a similar concentration.

\*This Product is intended for Laboratory Research use only.

## Binding of anti-CD45 antibody to human cell types

