

PERFORMANCE DATA SHEET

3325

Monoclonal anti-human CD51 (alpha v integrin)/Biotin *

mAb name/Clone: P2W7R

Isotype: Mouse IgG1 κ

Immunogen: Human ocular melanoma cell line (V+B2)

CATALOG#: 202-030

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD51 is an integrin alpha v chain which can associate with five different integrin beta chains to form various adhesion molecules. The alpha v beta 3 integrin complex binds vitronectin at the RGD sequence and also can bind to fibrinogen, von Willebrand factor, thrombospondin, fibronectin, osteopontin and collagen. Antibody P2W7 recognizes the alpha v molecule of 160 kd.

References: R.O. Hynes (1992) Cell 69: 11-25. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995) p. 1663-1664.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/thawing not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% Na₃N (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

PERFORMANCE: Five x 10⁵ cultured UM-SCC (squamous cell carcinoma) cells were harvested by trypsinization, washed and pre incubated 5 minutes with 20 μ l of 250 μ g/ml human IgG (to block nonspecific binding), after which they were incubated 45 minutes on ice with 80 μ l of anti-CD51/Biotin at 10 μ g/ml. Cells were washed twice and incubated with 2^o reagent Streptavidin/R-Phycoerythrin (Catalog #253-050), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.36 log₁₀ fluorescent units when compared to a Mouse IgG1/Biotin Isotype control (Catalog #278-030). Binding was blocked when cells were pre incubated 10 minutes with 20 μ l of 0.5 mg/ml anti-CD51 antibody (Catalog #202-020).

Binding of anti-CD51/Biotin +SA/PE to human UM-SCC cells



*Research use only. Not for use in Diagnostic procedures.