

PERFORMANCE DATA SHEET

1817

Monoclonal anti-human CD54(ICAM-1) Domain D2*

mAb name/Clone: 8.4A6

Isotype: Mouse IgG1

Immunogen: Human endothelial cells

CATALOG#: 206-020

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD54 (ICAM-1) mediates cell adhesion by binding to the integrins CD11a/CD18 (LFA-1) and to CD11b/CD18 (Mac-1). CD54 expression on resting peripheral blood leukocytes is weak but is upregulated on activated T and B lymphocytes and monocytes. Antibody 8.4A6 recognizes the D2 domain of the CD54 molecule of 90 kd. Antibody 8.4A6 inhibits CD54 binding to LFA-1.

References: A.R. Berendt, et al, (1992) Cell **68**: 71-81. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1548-1550. P.L. Reilly, et al, (1995) J Immunol **155**: 529-532.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10⁵ cultured **Raji** human tumor cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD54 antibody at a concentration of 5 µg/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.94 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog #278-010).

Binding of anti-CD54 mAb +GAM/FITC to human Raji cells



**This Product is intended for Laboratory Research use only.*