PERFORMANCE DATA SHEET

1850

Monoclonal anti-human CD55 (DAF)/FITC*

mAb name/Clone: 67 *Isotype:* Mouse IgG1k

Immunogen: Human monocytes

CATALOG#: 207-040 QUANTITY: 120 tests

VOLUME IN VIAL: 0.2ml

WORKING DILUTION: 1:50 (or use 1.6 μ l of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Human CD55 is a GPI anchored protein that can protect cells from complement mediated lysis. Antibody 67 recognizes the CD55 molecule of 70 kd.

References: D.G. Palmer, et al, (1985) Clin Exp Immunol 59: 529-538. Leukocyte Typing V (S.F. Schlossman, et al, eds.)

Oxford University Press, Oxford (1995) p. 1473-1474.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at **0.5 mg/ml** with a Fluorescein/IgG molar ratio of 3.7.

PERFORMANCE: Five x 10^5 cultured **HPB-MLT** cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD55/FITC at a dilution factor of **1:50** (10 µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.8** \log_{10} fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD55 antibody (Catalog #207-020).

*This Product is intended for Laboratory Research use only.

Isotype Control (red)

Blocked Control (blue)

anti-CD55/FITC (green)

102

10³

Binding of anti-CD55/FITC to human

HPB-MLT cells

100

101