

PERFORMANCE DATA SHEET

2348

Monoclonal anti-human CD57 (HNK-1)*

mAb name/Clone: NK-1

Isotype: Mouse IgMκ

Immunogen: Human PBL

CATALOG#: 209-020

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD57 originally called HNK-1 is a glycoprotein found on 15-20 percent of PBL's, including 60 percent of NK cells, and a subset of T cells (1). The immune regulation role of CD57 positive PBL's expressing high levels of CD8 is being investigated (2). Antibody NK-1 recognizes the CD57 molecule of about 110 kd.

References: 1) Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995) p. 1412-1414. 2) E.C.Y. Wang, et al, (1995) J Immunol 155: 5046-5056.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was purified to 95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin) using size exclusion chromatography.

PERFORMANCE: Five x 10⁵ cultured **Jurkat-4G** human tumor cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD57 antibody at 5 µg/ml. Cells were washed twice and incubated with 2⁰ reagent Goat anti-Mouse IgG/FITC (Catalog #232-011); after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.35 log₁₀ fluorescent units when compared to a Mouse IgM negative control (Catalog #290-010).

Binding of anti-CD57 mAb +GAM/FITC to human Jurkat-4G cells



**Research Use Only. Not for use in Diagnostic procedures.*