

PERFORMANCE DATA SHEET

1817

Monoclonal anti-human CD58(LFA-3)*

mAb name/Clone: TS2/9.1.4.3

Isotype: Mouse IgG1

Immunogen: Human cytolytic T cells

CATALOG#: 210-020

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD58 is a cell surface glycoprotein that functions as an adhesion molecule and mediates costimulatory activity through its ligand CD2, which is expressed on T and natural killer cells (1,2). CD58 has wide cellular distribution on both hemopoietic and non-hemopoietic cells with strong expression on macrophages and memory T cells. Antibody TS2 recognizes the CD58 glycoprotein of about 60-70 kd and will inhibit HLA-DR mediated T cell cytotoxicity.

References: (1) Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 414-415. (2) A.R. Arulanandam, et al, (1994) J Exp Med 180:1861-1871.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10⁵ cultured Nalm-6 cells were washed after which they were incubated 45 minutes on ice with 80 µl of anti-CD58 antibody at 10 µg/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.46 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog #278-010) at a similar concentration.

Binding of anti-CD58 antibody to human Nalm-6 cells



**This Product is intended for Laboratory Research use only.*