## PERFORMANCE DATA SHEET

181

## Monoclonal anti-human CD58(LFA-3)/Biotin\*

mAb name/Clone: TS2/9.1.4.3

Isotype: Mouse IgG1

Immunogen: Human cytolytic T cells

**CATALOG#: 210-030 QUANTITY: 100 μg** 

**CONCENTRATION: 1.0 mg/ml** 

**INFORMATION:** Human CD58 is a cell surface glycoprotein that functions as an adhesion molocule and mediates costimulatory activity through its ligand CD2, which is expressed on T and natural killer cells (1,2). CD58 has wide cellular distribution on both hemopoietic and non-hemopoietic cells with strong expression on macrophages and memory T cells. Antibody TS2 recognizes the CD58 glycoprotein of about 60-70 kd and will inhibit HLA-DR mediated T cell cytotoxicity.

**References:** (1) Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 414-415. (2) A.R. Arulanandam, et al, (1994) J Exp Med 180:1861-1871.

**STORAGE CONDITIONS:** *Store at 2 - 5°C*. Freeze/thawing not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

**PERFORMANCE:** Five x  $10^5$  cultured human **Nalm-6** cells were washed and incubated 45 minutes on ice with 80  $\mu$ l of anti-CD58/Biotin at **10**  $\mu$ g/ml. Cells were washed twice and incubated with  $2^0$  reagent Streptavidin/R-PE (Catalog #253-050) after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.1**  $\log_{10}$  fluorescent units when compared to a Mouse IgG1/Biotin negative control (Catalog # 278-030) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20  $\mu$ l of 0.5 mg/ml anti-CD58 antibody (Catalog #210-020).

\*This Product is intended for Laboratory Research use only.

## Binding of anti-CD58/Biotin to human Nalm-6 cells

