PERFORMANCE DATA SHEET 2140 Monoclonal anti-human CD59 (Protectin)/Biotin*



mAb name/Clone: BRA-10G *Isotype:* Mouse IgG1κ *Immunogen:* Human K562 tumor cells

CATALOG#: 211-030 QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD59 is a GPI anchored molecule that protects cells from complement-mediated lysis (1). Studies are progressing to determine the regulatory mechanism for CD59 expression (2). Antibody BRA-10G reacts with the 18-22 kd CD59 molecule.

References: (1) Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995) p. 1476-1477. (2) M.H. Holguin, et al, (1996) J Immunol 157: 1659-1668.

STORAGE CONDITIONS: Store at 2 - 5^oC. Freeze/thawing not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

PERFORMANCE: Five x 10^5 cultured human **HPB-MLT** cells were washed and pre incubated 5 minutes with 20 µl of 250 µg/ml human IgG (To block non specific binding) after which they were incubated 45 minutes on ice with 80 µl of anti-CD59/Biotin at **10 µg/ml**. Cells were washed twice and incubated with 2° reagent Streptavidin/R-PE (Catalog #253-050) after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.9** log₁₀ fluorescent units when compared to a Mouse IgG1/Biotin negative control (Catalog #278-030). Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD59 antibody (Catalog #211-020).

*Research use only. Not for use in Diagnostic procedures.



