PERFORMANCE DATA SHEET

3120

Monoclonal anti-human CD64 (FcγRI)*



mAb name/Clone: 10.1
Isotype: Mouse IgG1

Immunogen: Human rheumatoid synovial fluid cells/monocytes

CATALOG#: 216-820 (Preservative Free)

QUANTITY: 100 μg CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD64 is a high affinity receptor for monomeric human IgG1 and IgG3 which is expressed on macrophages, monocytes, and gamma interferon induced neutrophils. CD64 plays an important role in clearance of immune complexes and in antibody dependent cytoxicity. Antibody 10.1 recognizes the CD64 molecule of 72 kd from gene FcγRIA. Antibody 10.1 blocks binding of FcγRI to immunoglobin opsonized cells. *References:* G.J. Dougherty, et al, (1987) Eur J Immunol **17:** 1453-1459. Y. Jayaram, et al, (1989) Clin Exp Immunol **75:** 414-420. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 874-875. D.L. Durden, et al, (1995) J Immunol **154:** 4039-4047. L.L. Marnell, et al, (1995) J Immunol **155:** 2185-2193.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Open under aseptic conditions.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____


BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl. **No Preservative** added.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin. Product was 0.2μ sterile filtered and vialed under aseptic conditions.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10^5 cultured **U-937** cells were washed and incubated 45 minutes on ice with 80 μl of anti-CD64 antibody at **10** μg/ml. Cells were washed twice and incubated with 2^0 reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.83** log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog # 278-010).

Binding of anti-CD64 mAb +GAM/FITC to human U-937 cells



^{*} Research Use Only. Not for use in Diagnostic procedures.