

PERFORMANCE DATA SHEET

2323

Monoclonal anti-human CD66b (CEACAM8)***mAb name/Clone:** ANC4A4**Isotype:** Mouse IgG1 κ **Immunogen:** Recombinant soluble human CD66b**CATALOG#:** 220-020**QUANTITY:** 100 μ g**CONCENTRATION:** 1.0 mg/ml

INFORMATION: Human CD66b (CEACAM8, CGMb, NCA-95, CD67) is a 95kD member of the Carcinoembryonic Antigen (CEA) group. It is expressed on granulocytes and functions as an adhesion molecule. Similar to the CD66 a, c, and d isoforms, engagement of CD66b on neutrophils causes increased adhesion to endothelial cells (2). Crosslinking CD66b on neutrophils with antibodies triggers IL-8 release (3).

Antibody from clone ANC4A4 binds to CD66b on the cell surface of human PMN in Flow cytometry and works as an EIA capture reagent for recombinant CD66b utilizing either clone ANC1D5(Cat #217-030) or ANC3C5(Cat# 219-030) as a detection reagent.

References: 1) W.N. Khan, S. Hammarstrom, et al. (1992) *Genomics* **14**(2):384-90. 2) K.M. Skubitz, A.P. Skubitz, et al. (1996) *J Leuk Biol* **60**(1): 106-117. 3) A.K. Schroder, L Rink, et al. (2006) *Human Immunol* **67**(9): 676-682.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10⁵ Polymorphprep prepared human Granulocytes(PMN) per tube were pelleted and pre incubated 5 minutes with 20 μ l of human IgG at 300 μ g/ml (to reduce non specific binding) after which they were washed and incubated 45 minutes on ice with 80 μ l of anti-CD66b antibody(ANC4A4) at 5 μ g/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011) after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 0.91 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog #278-010) at a similar concentration. Binding was blocked when reagent was pre incubated with a 10-fold excess of recombinant CD66b.

***Research use only. Not for use in Diagnostic procedures.**

Binding of anti-CD66(ANC4A4) antibody +GAM/FITC to human PMN