

PERFORMANCE DATA SHEET

2635

Monoclonal anti-human CD66b (CEACAM8)*

mAb name/Clone: ANC4A4

Isotype: Mouse IgG1κ

Immunogen: Recombinant soluble human CD66b

CATALOG#: 220-820 (Preservative-free)

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD66b (CEACAM8, CGMb, NCA-95, CD67) is a 95kD member of the Carcinoembryonic Antigen (CEA) group. It is expressed on granulocytes and functions as an adhesion molecule. Similar to the CD66 a, c, and d isoforms, engagement of CD66b on neutrophils causes increased adhesion to endothelial cells (2). Crosslinking CD66b on neutrophils with antibodies triggers IL-8 release (3).

Antibody from clone ANC4A4 binds to CD66b on the cell surface of human PMN in Flow cytometry and works as an EIA capture reagent for recombinant CD66b utilizing either clone ANC1D5(Cat #217-030) or ANC3C5(Cat# 219-030) as a detection reagent.

References: 1) W.N. Khan, S. Hammarstrom, et al. (1992) *Genomics* **14**(2):384-90. 2) K.M. Skubitz, A.P. Skubitz, et al. (1996) *J Leuk Biol* **60**(1): 106-117. 3) A.K. Schroder, L. Rink, et al. (2006) *Human Immunol* **67**(9): 676-682.

STORAGE CONDITIONS: Store at 2 - 5°C. Open under aseptic conditions. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin. Product was 0.2 µm filtered and vialled under aseptic conditions.

PERFORMANCE: Five x 10⁵ Polymorphprep prepared human Granulocytes(PMN) per tube were pelleted and pre incubated 5 minutes with 20µl of human IgG at 300µg/ml (to reduce non specific binding) after which they were washed and incubated 45 minutes on ice with 80 µl of anti-CD66b antibody(ANC4A4) at 5 µg/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011) after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 0.91 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog #278-010). Binding was blocked when reagent was pre incubated with a 10-fold excess of recombinant CD66b.

Binding of anti-CD66(ANC4A4) antibody +GAM/FITC to human PMN



* Research Use Only. Not for use in Diagnostic procedures.