

PERFORMANCE DATA SHEET

3328

Monoclonal anti-human CD69(AIM)*

mAb name/Clone: HP-4B3

Isotype: Mouse IgG2a

Immunogen: IL-2 activated human NK cells

CATALOG#: 221-820 (Preservative Free)

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD69 is expressed by platelets and mature thymocytes and appears rapidly after activation on T cells, B cells and NK cells. This pattern of cellular CD69 expression has suggested a lymphocyte activation role. Functional studies with antibodies to CD69 also support an activation role for CD69. Antibody HP-4B3 recognizes the dimeric CD69 molecule of 60 kd.

References: S.F. Ziegler, et al, Stem Cells (1994) **12**: 456-465. R. Testi, et al, Immunol Today (1994) **15**: 479-483. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1123-1132. D. Vanhecke, et al, J Immunol (1995) **155**: 1862-1872.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. *Open under aseptic conditions.*

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____


BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl. **No preservative added.**

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin. Product was 0.2µ sterile filtered and viald under aseptic conditions.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10⁵ cultured **HEL** cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD69 antibody at **10 µg/ml**. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.40 log₁₀** fluorescent units when compared to a Mouse IgG2a negative control (Catalog #281-010).

Binding of anti-CD69 mAb +GAM/FITC to human HEL92 cells



**Research use only. Not for use in Diagnostic procedures.*