

*Monoclonal anti-human CD71 (Transferrin receptor)**

mAb name/Clone: DF1513

Isotype: Mouse IgG1

Immunogen: Human PBL

CATALOG#: 223-825 (Preservative free)

QUANTITY: 2 mgs

CONCENTRATION: 10.0 mg/ml

INFORMATION: Human CD71 is a type II transmembrane glycoprotein found on activated leukocytes. CD71 binds transferrin and mediates uptake of iron through endocytosis and recycling. Antibody DF1513 recognizes the CD71 molecule of about 95 kd.

References: 1)Leukocyte Typing IV (W. Knapp, et al, eds.) Oxford University Press, Oxford, (1989) p. 455-460. 2)Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 524-527.

STORAGE CONDITIONS: *Store at 2 - 5°C. Open under aseptic conditions.* Freeze/Thawing is not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM Sodium Chloride.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10⁵ cultured **HPB-MLT** cells were washed and preincubated 5 minutes with 20 µl of 250 µg/ml human IgG (To block non specific binding) after which they were incubated 45 minutes on ice with 80 µl of anti-CD71 antibody at **5 µg/ml**. Cells were washed twice and incubated with 50 µl of 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.81 log₁₀** fluorescent units when compared to a Mouse IgG1 negative control (Catalog # 278-010).

** Research Use Only. Not for use in Diagnostic procedures.*

Binding of anti-CD71 mAb +GAM/FITC to human HPB-MLT cells

