PERFORMANCE DATA SHEET 1817 Monoclonal anti-human CD74/FITC*

mAb name/Clone: M-B741 *Isotype:* Mouse IgG2a *Immunogen:* RPMI 1788 Cell Line

 CATALOG#: 226-040
 VOLUME IN VIAL: 0.2 ml

 QUANTITY: 120 tests
 VOLUME IN VIAL: 0.2 ml

 WORKING DILUTION: 1:50 (1.6μl conc stock per 5 x 10⁵ cell test)

INFORMATION: Human CD74 is expressed intracellularly and on the cell surface of MHC Class II positive cells such as B cells, monocytes, macrophages, Langerhans cells and dendritic cells. CD74 has several important functions relating to antigen presentation by MHC Class II molecules. Antibody M-B741 recognizes the CD74 molecule of about 33 kd.

References: Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 568-574. P. Romagnoli & R.N. Germain, (1995) J Exp Med **182:** 2027-2036.

STORAGE CONDITIONS: Store at 2 - 5^oC. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at approximately **0.5 mg/ml** with a Fluorescein:Ab molar ratio of 7.7. **Binding of anti-CD74/FT**

PERFORMANCE: Five x 10^5 cultured **Raji** cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD74/FITC at a **1:50** dilution. Cells were then washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.8** log₁₀ fluorescent units when compared to a Mouse IgG2a/FITC negative control (Catalog #281-040) at a similar concentration. This shift was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD74 antibody (Catalog #226-020).

*This Product is intended for Laboratory Research use only.

Binding of anti-CD74/FITC to human Raji Cells

