

**PERFORMANCE DATA SHEET**  
2448

**Monoclonal anti-human CD79a(mb1)/FITC\***

**mAb name/Clone:** ZL7-4  
**Isotype:** Mouse IgG1  
**Immunogen:** Daudi IgM complex

**CATALOG#:** 235-040

**QUANTITY:** 120 tests

**VOLUME IN VIAL:** 0.2ml

**WORKING DILUTION:** 1:50 (or use 1.6µl of concentrated stock per 5 x 10<sup>5</sup>-cell test)

**INFORMATION:** Human CD79a is a 22kd glycoprotein which associates with CD79b to form a functional heterodimer that interacts with membrane immunoglobulin (mIg). The mIgM component binds antigen and the heterodimer transduces signals. Antibody ZL7-4 recognizes the extracellular part of CD79a.

**References:** 1) L. Zhang, M.J. Glennie, et al. (1995) *Therapeutic Immunol.* 2: 191-202. 2) Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 180-185. 3) Chu PG, Arber DA. (2001) *Appl Immunohistochem Molecul Morphol* 9: 97-106.

**STORAGE CONDITIONS:** Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: \_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at approximately 0.5 mg/ml with a Fluorescein/IgG molar ratio of 5.6.

**PERFORMANCE:** Five x 10<sup>5</sup> cultured human Daudi cells were pre incubated 5 minutes with 20 µl of 250 µg/ml human Ig (to block non specific binding) after which they were incubated 45 minutes on ice with 80 µl of product at a 1:50 dilution (10 µg/ml). They were then washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.65 log<sub>10</sub> fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040). Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD79a antibody (Catalog #235-020).

\* *Research Use Only. Not for use in Diagnostic procedures.*

**Binding of anti-CD79a/FITC to human Daudi cells**

