

PERFORMANCE DATA SHEET

2036

Monoclonal anti-human IL-2*

mAb name/Clone: ANC7F7

Isotype: Mouse IgG1 κ

Immunogen: Recombinant human IL-2

CATALOG#: 245-020

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: The cytokine human interleukin-2 (IL-2) is a 15 kd protein expressed upon activation by peripheral TH1 cells, medullary thymocytes, and a subset of large granular lymphocytes. Secreted IL-2 N7-48A stimulates proliferation and lymphokine production by T cells, B cells and NK cells. Antibody ANC7F7 binds to soluble and intracellular IL-2 in FACS and EIA.

References: 1. S.L. Swain (1991) *Curr Opin Immunol* 3:304-310. 2. A.K. Abbas, et al, (1996) *Nature* 383: 787-793. 3. A. Rebollo, et al, (1996) *Advan Immunol* 63: 127-196.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

PERFORMANCE: Ficoll prepared human peripheral blood mononuclear cells were stimulated 4 hours in the presence of 50 ng/ml PMA, 2 μ M ionomycin and 5 μ M monensin. Cells were then harvested, washed and fixed for 30 min with 2% buffered formaldehyde after which they were washed two times. Fixed cells were permeabilized 10 minutes in a buffer containing 0.3% Saponin. Subsequent reagent incubations and washes were done using this buffer. Five x 10⁵ cells per tube were pre incubated 5 minutes with 20 μ l of 250 μ g/ml human IgG (to block non specific binding), after which they were incubated 45 minutes on ice with 80 μ l of anti-IL-2 antibody at 5 μ g/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. A net 40% sub population of the cells stained positive with a mean shift of 1.07 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog # 278-010) at a similar concentration.

Binding of anti-IL-2 mAb +GAM/FITC to stimulated fixed permeabilized human PBMC



* **Research Use Only. Not for use in Diagnostic procedures.**