## PERFORMANCE DATA SHEET

1818

## Monoclonal anti-mouse CD8 alpha (Lyt 2)\*

*mAb name/Clone:* 53-6.72 *Isotype:* Rat IgG2ak

Immunogen: murine thymocytes/splenocytes

**CATALOG#: 260-020 QUANTITY: 100 μg** 

**CONCENTRATION: 1.0 mg/ml** 

**INFORMATION:** Antibody 53-6.72 binds to the alpha subunit of murine CD8 for all mouse strains tested. It is a useful tool for assessing murine CD8α containing recombinant proteins. **References:** 1) Ledbetter, J.A., et al. (1979) Immunol. Rev. 47: 63-90.

**STORAGE CONDITIONS:** *Store at 2 - 5<sup>o</sup>C*. Freeze/Thawing is not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_


**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

**PRODUCTION:** Antibody was Protein G purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

**PERFORMANCE:** Reagent was tested for binding to ACK lysed murine splenocytes in FACS. Five x 10<sup>5</sup> **splenocytes** per tube were washed and pre incubated with 20μl of 300μg/ml murine IgG (to reduce non specific

binding) after which they were incubated 45 minutes on ice with 80  $\mu$ l of anti-CD8 antibody diluted to  $10~\mu g/ml$ . Cells were washed twice and incubated with  $2^{\rm O}$  reagent Goat anti-Rat IgG/FITC, after which they were washed three times, fixed and analyzed by FACS. An 8.8~% sub population of the cells stained positive with a mean shift of  $1.6~\log_{10}$  fluorescent units when compared to background.

Binding of anti- Mouse CD8 mAb + GAR/FITC to Balb-c splenocytes



\*This Product is intended for Laboratory Research use only.