

PERFORMANCE DATA SHEET

1818

Monoclonal anti-human CD278(ICOS)/FITC*

mAb name/Clone: ANC6C6

Isotype: Mouse IgG1 κ

Immunogen: Human HPB-MLT cells, human ICOS-muIg fusion protein

CATALOG#: 265-050

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2 ml

WORKING DILUTION: 1:50 (or use 1.6 μ l of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: The inducible costimulator (ICOS, T cell activation molecule H4) is similar to human CD28 (24% homology), and plays an analogous role in the T cell activation process. Secondary signaling through CD28 or ICOS results in discrete cytokine secretion profiles by the activated T cells.¹ Signaling by either molecule is effectively down regulated by CD152 (CTLA-4) engagement.² Human GL50 is a member of the B7 family sharing ~20% homology with CD80 (B7-1) and CD86 (B7-2), and has been shown to be the ligand for ICOS.³ Two RNA splice variants exist for this molecule, differing only in the cytoplasmic domain.⁴

Antibody ANC6C6 binds to recombinant ICOS-muIg in EIA, and to the surface of HPB-MLT cells in Flow cytometry. Additionally, it blocks binding of recombinant GL50-muIg to HPB-MLT cells.

References: 1) Beier, K.C., R.A. Kroczek, et al. 2000, *Eur J Immunol.* **30**(12):3707-3717. 2) Riley, J.L., C.H. June, et al. 2001, *J. Immunol.* **166**: 4943-4948. 3) Ling, V., M. Collins, et al. 2000, *J. Immunol.* **164**: 1653-1657. 4) Ling, V., M. Collins, et al. 2001, *J. Immunol.* **166**: 7300-7308.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at 0.5 mg/ml with a Fluorescein/IgG molar ratio of 6.4.

PERFORMANCE: Five x 10⁵ cultured human HPB-MLT cells per tube were washed and incubated 45 minutes on ice with 80 μ l of product at a dilution factor of 1:50 (10 μ g/ml). They were then washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 0.47 log₁₀ fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 μ l of 0.5 mg/ml anti-ICOS antibody (Catalog #265-020).

**This Product is intended for Laboratory Research use only.*

R-Phycoerythrin (R-PE) is covered under patents: U.S. 4,520,110; European 76,695 and Canadian 1,179,942.

Binding of anti-CD278(ICOS)/FITC to human HPB-MLT Cells

