

PERFORMANCE DATA SHEET

2916

Monoclonal anti-human CD257(BAFF, Tall-1)/R-PE*

mAb name/Clone: ANC2H3

Isotype: Mouse IgG1k

Immunogen: Recombinant CD257(BAFF)

CATALOG#: 266-050

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2 ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Human CD257 (B cell activating factor, BAFF,TALL-1, Blys, THANK) and CD256 (APRIL, a proliferation inducing ligand) are both type II molecules belonging to the TNF superfamily (TNFSFL #13b and 13 respectively). They are expressed by non-B cells, and are down regulated by mitogenic stimulation(2). BAFF and APRIL bind to at least two receptors: CD267 (TACI, transmembrane activator and CAML-interactor) and CD269 (BCMA, B cell maturation antigen), both of which are restricted to B cells(3,4). Ligation of these receptors with recombinant BAFF dramatically increases IgM production by peripheral blood B cells(1). A third receptor CD268 (BAFFR) is specific for BAFF has also been described(5). BAFF and BAFFR knockout mice have a reduced numbers of mature B cells in the periphery, however TACI and BCMA knockouts do not share this phenotype, suggesting that BAFF-R may the primary receptor for BAFF in mice(8,9,10). Cell surface BAFF can be proteolytically cleaved to form a soluble trimeric molecule(2). Levels of soluble BAFF correspond with levels of autoantibodies in Sjogren's Syndrome(11). Clone ANC2H3 blocks binding of recombinant human CD257(BAFF) to receptors on Raji cells in Flow Cytometry.

References: 1) Schneider P., J. Tschopp, et al. *J. Exp. Med.* 1999, 189(11):1747-1756. 2) Shu, H.B., H. Johnson, W.H. Hui. *J Leukoc Biol* 1999, 65:680-683. 3) Marsters, S.A., A. Ashkenazi, et al. 2000, *Curr Biol* 10:785-788.

4) Xia, X., H. Hsu, et al. 2000, *J Exp Med.* 192(1): 137-143. 5) Thompson J.S., C. Ambrose, et al. *Science* 2001, 293: 2108-2111. 6) Roschke, V, T.S. Migone, et al. *J Immunol.* 2002, 169: 4314-4321. 7) MacLennan, C.M., C.G. Vinuesa, 2002, *Immunity* 17:235-238. 8) B. Schiemann, et al, (2001) *Science* 293: 2111-2114. 9) S.M. Harless, et al, (2001) *Curr Biol* 11: 1988-1989. 10) *Mol Cell Biol* (2001) 21: 4067-4074. 11) X. Mariette, et al, (2003) *Ann Rheum Dis* 62: 168-171.

STORAGE CONDITIONS: Store at 2 - 5°C. Do not freeze! Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at 0.5 mg/ml with an A₅₆₅/A₂₈₀ ratio of 3.04.

PERFORMANCE: Five x 10⁵ recombinant CD257-muCD8-coated CML Latex beads (inquire!) per tube were washed and incubated 45 minutes on ice with 80 µl of anti-CD257/PE a dilution factor of 1:50 (10 µg/ml). They were washed three times, fixed and analyzed by FACS. Beads stained positive with a mean shift of 3.6 log₁₀ fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050) at a similar concentration. Binding was blocked when beads were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD257 antibody (Catalog #266-020).

* **Research Use Only. Not for use in Diagnostic procedures.**

Binding of anti-CD257(BAFF)/PE to Recombinant CD257-coated Beads

