

PERFORMANCE DATA SHEET

1818

anti-human LT β R (Lymphotoxin beta receptor)/Biotin*

mAb name/Clone: ANCLTR2/9E2

Isotype: Mouse IgG1 κ

Immunogen: Recombinant human LT β R

CATALOG#: 267-030

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human Lymphotoxin beta receptor (LT β R, TNFSF3) is a member of the TNF receptor superfamily with similarity to CD120a(TNFR1) and CD120b(TNFR2). It is expressed mainly on non lymphoid tissues(1) and appears to have an important role in secondary lymphoid organ development(5). Suppression of LT β R signaling can alleviate autoimmunity(2) or exasperate mycobacterial infection(6). Its ligands include Lt $\alpha\beta_2$ and LIGHT. In mice, LT β R signaling is important for development and function of HEV(2), Dendritic cells(3), and Mast cells(4).

Antibody ANCLTR2 binds to human LT β R present on U-937 cell surface, and to recombinant LT β R- μ Ig in EIA.

REFERENCES: 1) T W Mak, M E Saunders eds. *The Immune Response* 2006 Elsevier Academic Press p 499.

2) J L Browning, R.A. Fava, et al. *Immunity* (2005) **23**(5): 539-50. 3) Y Wang, Y Fu, et al. *J Immunol* (2005) **175**: 6997-7002. 4) P Stopfer, T Hehlhans, et al. *J Immunol* (2004) **172**: 7459-7465. 5) Gommermann J, JL Browning, et al, *Nat Rev Immunol* (2003) **3**:642-655. 6) Spahn TW, T Kucharzik, et al. *Infection and Immunity* (2005) **73**(11): 7077-7088.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/thawing not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

PERFORMANCE: Five x 10⁵ cultured human U-937 cells were washed and pre incubated 5 minutes with 20 μ l of 250 μ g/ml human IgG (to block non specific binding) after which they were incubated 45 minutes on ice with 80 μ l of anti-LT β R/Biotin at 10 μ g/ml. Cells were washed twice and incubated with 2^o reagent Streptavidin/R-PE (Catalog #253-050) after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 0.64 log₁₀ fluorescent units when compared to a Mouse IgG1/Biotin negative control (Catalog #278-030) at a similar concentration. Binding was blocked when cells were pre blocked with 20 μ l of 0.5 mg/ml unlabeled anti-LT β R antibody (Catalog #267-020).

**This Product is intended for Laboratory Research use only.*

Binding of anti-LT β R/Biotin + SA/PE to human U-937 cells

