

PERFORMANCE DATA SHEET

3326

Monoclonal anti-human CD357 (GITR, AITR)*

mAb name/Clone: ANC7D6-1B7

Isotype: Mouse IgMk

Immunogen: Recombinant human GITR, PHA stimulated human PBL

CATALOG#: 268-020

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Glucocorticoid induced Tumor Necrosis Factor receptor family related gene (GITR, AITR) is a type 1 228 amino acid transmembrane protein that has been designated TNFRSF18. GITR is expressed mainly by CD4+ CD25+ T lymphocytes. Its engagement by antibodies or its ligand (GITRL) induces a costimulatory signal which can break T cell suppression (2). GITR has been implicated as a regulatory mechanism in anti-parasitic immune responses (4), and is involved with Macrophages activation in atherosclerotic plaque formation (3). Clone ANC7D6-1E7 binds to human GITR in EIA and on stimulated cell surface in FACS. Pre incubation of the receptor with an excess of this mAb will block binding of recombinant GITR Ligand.

References: 1) Gurney AL, et al. (1999) *Curr Biol* **9**(4): 215-218. 2) Ji HB, et al. (2004) *J Immunol* **172**(10): 5823-5827. 3) Kim WJ, WH Lee, et al. (2006) *Immunology* **119**(3): 44-9. 4) Furze RC, Selkirk ME, et al. (2006) *Microbes Infect* **8**(12-13): 2803-10. 5) Kohm AP, SD Miller, et al. (2004) *J Immunol* **172**: 4686-4690. 6) McHugh, R.S. et al. (2002) *Immunity* **16**:311. 7) Shin, H.H. et al. (2002) *FEBS Lett.* **514**:275. 8). Shimizu, J. et al. (2002) *Nature Immunol.* **3**:135.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was purified to 95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin) using size exclusion chromatography.

PERFORMANCE: Ficoll prepared human peripheral blood mononuclear cells (PBMC) were stimulated 5 days in culture with PHA. Five x 10⁵ cells per tube were washed and pre incubated 5 minutes with 20 µl of 250 ug/ml human IgG (to block non specific binding) after which they were incubated 45 minutes on ice with 80 µl of anti-GITR antibody at 5 µg/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011; This reagent cross reacts well with Mouse IgM), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 0.78 log₁₀ fluorescent units when compared to a Mouse IgM negative control (Catalog #290-010).

Binding of anti-CD357(GITR) mAb +GAM/FITC to stimulated human PBL



* **Research Use Only. Not for use in Diagnostic procedures.**