## PERFORMANCE DATA SHEET 2528 Monoclonal anti-human CD270(HVEM)/Biotin\*



*mAb name/Clone:* ANC3B7 *Isotype:* Mouse IgG2aκ *Immunogen:* Recombinant human CD270-muIg fusion protein

## CATALOG#: 270-030 CONCENTRATION: 1.0 mg/ml

## QUANTITY: 100 µg

Human CD270 (Herpes Virus Entry Mediator, TNFRSF14, ATAR, LIGHTR, TR2) is a 36kd type I membrane protein that has been found on a broad range of lymphoid tissues. Its ligands include the TNF Ligand family members Ltα and LIGHT, and the CD28/Ig Superfamily member BTLA(2). Antibody ANC3B7 is reactive with recombinant CD270 in EIA, and binds to CD270 present on human U-937 cell surface in Flow Cytometry.

**REFERENCES:** (1) Hsu H, et al. (1997) *J Biol Chem* **272**:13471. (2) Gonzalez LC, et al. (2005) *PNAS* **102**(4): 1116-1121.

**STORAGE CONDITIONS:** Store at 2 - 5<sup>o</sup>C. Freeze/thawing not recommended.


**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_


**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin.

Unconjugated Biotin was removed from conjugate using a desalting column.

**PERFORMANCE:** Five x  $10^5$  cultured human **U-937** cells per tube were washed and pre incubated 5 minutes with 20 µl human IgG at 250 µg/ml (to block non specific binding) after which they were incubated 45 minutes on ice with 80 µl of anti-CD270/Biotin at a concentration of **10 µg/ml**. Cells were washed twice and incubated with 2<sup>o</sup> reagent Streptaviden/R-PE (Catalog #253-050), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.47** log<sub>10</sub> fluorescent units when compared to a Mouse IgG2a/Biotin negative control (Catalog #281-030). Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD270 antibody (Catalog #270-020).





\*Research use only. Not for use in Diagnostic procedures.

Ancell Corporation P.O. Box 87 Bayport, MN 55003-0087 USA Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940