

PERFORMANCE DATA SHEET

1906

Monoclonal anti-human CD79b(BCR-Ig β)/Biotin*

mAb Name/Clone: SN8/3A2-2E7

Isotype: Mouse IgG1 κ

Immunogen: Membrane preparation of human B prolymphocytic leukemia cells

CATALOG#: 301-030

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD79b is the B cell antigen receptor Ig β chain (BCR-Ig β) which associates with B cell antigen receptor Ig α chain to form a functional heterodimer that interacts with membrane immunoglobulin (mIg). The mIgM component binds antigen and the Ig α /Ig β heterodimer transduces signals. Antibody SN8 recognizes the extracellular part of the BCR complex β chain (CD79b) which has a molecular weight of approximately 38 kd

References: M. Okazaki, et al, (1993) *Blood* **81**: 84-94. S. Vasile, et al, (1994) *Mol Immunol* **31**: 419-427. A.L. DeFranco, et al, (1994) *Chem Immunol* **59**: 156-172. *Leukocyte Typing V* (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 667-675, 677-681. Chu PG, Arber DA. (2001) *Appl Immunohistochem Molecul Morphol* **9**: 97-106.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/thawing not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

PERFORMANCE: Five x 10⁵ cultured Daudi cells were washed and incubated 45 minutes on ice with 80 μ l of anti-CD79b/Biotin at 10 μ g/ml. Cells were then washed twice and incubated 45 minutes with 2^o reagent Streptavidin/R-Phycoerythrin (Catalog #253-050), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 2.09 fluorescent units when compared to a Mouse IgG1/Biotin negative control (Catalog #278-030) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 μ l of 0.5 mg/ml anti-CD79b antibody (Catalog #301-020).

**This Product is intended for Laboratory Research use only.*

Binding of anti-CD79b/Biotin +SA/PE to human Daudi cells

