

PERFORMANCE DATA SHEET

1818

*Monoclonal anti-human CD84/FITC**

mAb name/Clone: 152.1D5

Isotype: Mouse IgG1

Immunogen: Human MLR stimulated lymphocytes

CATALOG#: 305-040

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5×10^5 -cell test)

INFORMATION: Human CD84 is a cell surface glycoprotein found on mature B lymphocytes, monocytes, and a subset of T lymphocytes. Antibody 152.1D5 recognizes the CD84 molecule of about 74 kd (1,2).

References: (1) Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995) p. 699-700, 1099, 1102. (2) Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 193-195.

STORAGE CONDITIONS: *Store at 2 - 5°C.* Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN_3 (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at **0.5 mg/ml** with a Fluorescein/IgG molar ratio of 9.1.

PERFORMANCE: Five $\times 10^5$ cultured **Raji** cells were incubated 45 minutes on ice with 80 µl of anti-CD84/FITC at a **1:50** dilution (10µg/ml). They were then washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.81** \log_{10} fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD84 antibody (Catalog #305-020).

**This Product is intended for Laboratory Research use only.*

Binding of anti-CD84/FITC to human cell lines

