

PERFORMANCE DATA SHEET

2335

Monoclonal anti-human CD94 (Kp43)/FITC*

mAb name/Clone: HP-3D9

Isotype: Mouse IgG1

Immunogen: Cultured human NK cells

CATALOG#: 315-040

QUANTITY: 120 tests

VOLUME IN VIAL: 100 µl

WORKING DILUTION: 1:50 (or use 1.6 µl of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Human CD94 expression is restricted to natural killer (NK) cells and a small subset of T lymphocytes. CD94 is a type II membrane glycoprotein of the c-type lectin superfamily and essentially lacks a cytoplasmic tail. Antibody HP-3D9 recognizes the CD94 molecule of 70 kd. Antibody HP-3D9 inhibits IL-2 dependent proliferation of NK cells.

References: J. Aramburu, et al, (1990) J Immunol **144**: 3238-3247. J. Aramburu, et al, J Immunol (1991) **147**: 714-721. A. Moretta, et al, J Exp Med (1994) **180**: 545-555. C. Chang, et al, Eur J Immunol (1995) **25**: 2433-2437. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p.1437-1439.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at **0.5 mg/ml** with a Fluorescein/IgG molar ratio of 7.2.

PERFORMANCE: Ficoll prepared **human peripheral white blood cells** were stimulated by incubating 1 day at 5 x 10⁶ cells/ml in RPMI 10% FBS media including 5 µg/ml Phytohemagglutinin-P. Five x 10⁵ cells per tube were then washed and incubated 45 minutes on ice with 80 µl of anti-CD94/FITC at a **1:50** dilution (10 µg/ml). Cells were washed three times, fixed and analyzed by FACS. A 27% sub population of the cells stained positive with a mean shift of **1.22** log₁₀ fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD94 antibody (Catalog #315-020).

Binding of anti-CD94/FITC to stimulated human PBMC



**Research Use Only. Not for use in Diagnostic procedures.*