## PERFORMANCE DATA SHEET

1212

## Ancell

## Monoclonal anti-human CD95 (APO-1/FAS)/R-PE\*

mAb name/Clone: ANC95.1/5E2

*Isotype:* Mouse IgG1

Immunogen: Human soluble FAS with the transmembrane region deleted

CATALOG#: 316-050
QUANTITY: 120 tests

VOLUME IN VIAL: 0.2 ml
WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5 x 10<sup>5</sup>-cell test)

**INFORMATION:** Human CD95 (APO-1/FAS) is a type I cell surface glycoprotein that is strongly upregulated on activated T cells, B cells, NK cells and thymocytes. CD95 plays an important role in programmed cell death or apoptosis. Apoptosis appears to be a mechanism for regulating the immune response.

References: Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1142-1148. S. Nagata & P. Golstein (1995) Science 267: 1449-1456. S. Nagata & T. Suda (1995) Immunol Today 16: 39-43. D.H. Lynch, F. Ramsdell & M.R. Alderson (1995) Immunol Today 16: 569-574.

STORAGE CONDITIONS: Store at 2 - 5°C. Do not freeze! Protect from light.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at **0.5 mg/ml** with an  $A_{565}/A_{280}$  ratio of 2.96.

**PERFORMANCE:** Human **Raji** cells were incubated 45 minutes on ice with 80  $\mu$ l of anti-CD95/R-PE at a dilution factor of **1:50** (10  $\mu$ g/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **2.01**  $\log_{10}$  fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050). Binding was blocked when cells were pre incubated 10 minutes with 20  $\mu$ l of 0.5 mg/ml anti-CD95 antibody (Catalog #316-020).

\* Research Use Only. Not for use in Diagnostic procedures.

## Binding of anti-CD95/R-PE to human cell types

