

**PERFORMANCE DATA SHEET**

3248

**Monoclonal anti-human CD106(VCAM-1)/Biotin\*****mAb name/Clone:** 1.G11B1**Isotype:** Mouse IgG1**Immunogen:** Human endothelial cells**CATALOG#:** 327-030**QUANTITY:** 100 µg**CONCENTRATION:** 1.0 mg/ml

**INFORMATION:** Human CD106 is an endothelial adhesion molecule that binds to  $\alpha_4\beta_1$  &  $\alpha_4\beta_7$  integrins (VLA-4) and promotes adhesion of lymphocytes, monocytes, eosinophils and basophils. CD106 is also found on follicular dendritic cells, interdigitating reticulum cells and Kupffer cells. Antibody 1.G11B1 recognizes the VCAM-1 molecule (CD106) and blocks leukocyte adhesion.

**References:** M.H. Thornhill, D.O. Haskard, (1991) J Immunol **146**: 592-598. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1764-1767. M. Nagata, et al, (1995) J Immunol **155**: 2194-2202. H.E. Chuluyan, et al, (1995) J Immunol **155**: 3135-314.


**STORAGE CONDITIONS:** *Store at 2 - 5°C.* Freeze/thawing not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: \_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% Na<sub>3</sub> (as a preservative).

**PRODUCTION:** Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

**PERFORMANCE:** Cultured **human umbilical cord vein endothelial cells** were stimulated 4 hours with 10 ng/ml Phorbol 12-Myristate 13-Acetate. Five x 10<sup>5</sup> cells per tube were then washed and incubated 45 minutes on ice with 80 µl of anti-CD106/Biotin at **10 µg/ml**. Cells were washed twice and incubated with 2<sup>o</sup> reagent, Streptavidin/R-Phycoerythrin (Catalog #253-050), after which they were washed three times, fixed and analyzed by FACS. A 34% sub population of the cells stained positive with a mean shift of **1.02 log<sub>10</sub>** fluorescent units when compared to a Mouse IgG1/Biotin negative control (Catalog #278-030) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD106 antibody (Catalog #327-020).

**Binding of anti-CD106/Biotin + SA/PE to Stimulated HUVEC**

*\*This Product is intended for Laboratory Research use only.*