

PERFORMANCE DATA SHEET

1947

Monoclonal anti-human CD106(VCAM-1)/FITC***mAb name/Clone:** 1.G11B1**Isotype:** Mouse IgG1**Immunogen:** Human endothelial cells**CATALOG#:** 327-040**QUANTITY:** 120 tests**VOLUME IN VIAL:** 0.2 ml**WORKING DILUTION:** 1:50 (or use 1.6µl of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Human CD106 is an endothelial adhesion molecule that binds to $\alpha_4\beta_1$ & $\alpha_4\beta_7$ integrins (VLA-4) and promotes adhesion of lymphocytes, monocytes, eosinophils and basophils. CD106 is also found on follicular dendritic cells, interdigitating reticulum cells and Kupffer cells. Antibody 1.G11B1 recognizes the VCAM-1 molecule (CD106) and blocks leukocyte adhesion.

References: M.H. Thornhill, D.O. Haskard, (1991) J Immunol **146**: 592-598. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1764-1767. M. Nagata, et al, (1995) J Immunol **155**: 2194-2202. H.E. Chuluyan, et al, (1995) J Immunol **155**: 3135-314.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% Na₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at **0.5 mg/ml** with a Fluorescein/Antibody molar ratio of 8.2.

PERFORMANCE: Cultured **human umbilical cord vein endothelial cells** were stimulated 4 hours with 10ng/ml Phorbol 12-Myristate 13-Acetate. Five x 10⁵ cells per tube were then washed and incubated 45 minutes on ice with 80 µl of anti-CD106/FITC at a **1:50** dilution (10 µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.6** log₁₀ fluorescent units when compared to a Mouse IgG1/FITC negative control (Catalog #278-040) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD106 antibody (Catalog #327-020).

***For Research Use Only. Not For Use In Diagnostic Procedures.**

Binding of anti-CD106/FITC to stimulated HUVEC

