

PERFORMANCE DATA SHEET

3248

Monoclonal anti-human CD106(VCAM-1)/R-PE*

mAb name/Clone: 1.G11B1

Isotype: Mouse IgG1

Immunogen: Human endothelial cells

CATALOG#: 327-050

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2 ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated reagent per 5x10⁵-cell test)

INFORMATION: Human CD106 is an endothelial adhesion molecule that binds to α₄β₁ & α₄β₇ integrins (VLA-4) and promotes adhesion of lymphocytes, monocytes, eosinophils and basophils. CD106 is also found on follicular dendritic cells, interdigitating reticulum cells and Kupffer cells. Antibody 1.G11B1 recognizes the VCAM-1 molecule (CD106) and blocks leukocyte adhesion.

References: M.H. Thornhill, D.O. Haskard, (1991) J Immunol **146**: 592-598. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1764-1767. M. Nagata, et al, (1995) J Immunol **155**: 2194-2202. H.E. Chuluyan, et al, (1995) J Immunol **155**: 3135-314.

STORAGE CONDITIONS: *Store at 2 - 5°C. Do not freeze! Protect from light.*

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at **0.5 mg/ml** with an A₅₆₅/A₂₈₀ ratio of 3.53.

PERFORMANCE: Cultured **human umbilical cord vein endothelial cells** were stimulated 4 hours with 10 ng/ml Phorbol 12-Myristate 13-Acetate. Five x 10⁵ cells per tube were harvested, washed and incubated 45 minutes on ice with 80 µl of anti-CD106/R-PE at a **1:50** dilution (10µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.36 log₁₀** fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog # 278-050) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD106 antibody (Catalog #327-020).

*** Research Use Only. Not for use in Diagnostic procedures.**

Binding of anti-CD106/PE to stimulated HUVEC

