

PERFORMANCE DATA SHEET

2546

Monoclonal anti-human Hif-1 α *

mAb name/Clone: ANC10G3

Isotype: Mouse IgG1k

Immunogen: Recombinant human Hif1 α

CATALOG#: 335-020

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: Hypoxia Inducible Factor 1 alpha (Hif1 α) is a transcription factor which accumulates in the nucleus of cells in most tissues which are subjected to hypoxic conditions. There it stabilizes by associating with Hif1 beta (ARNT). Clone ANC10G3 binds to recombinant human Hif1 α in EIA, as well as native Hif1 α present in nuclei of fixed permeabilized Nalm-6 cells cultured overnight with CoCl₂ to simulate hypoxia. It interferes with binding of anti-Hif1 α clone 546-16, suggesting that their epitopes overlap.

Hif1 References: 1) GL Wang, GL Song, et al. (1993) *PNAS* **90**: 4304.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin. Product was 0.2 μ m filtered and vialled under aseptic conditions.

PERFORMANCE: Antibody specific activity was tested in FACS. Human Nalm-6 cells were cultured overnight in the presence of 200 μ M CoCl₂ to simulate hypoxia, after which they were fixed and permeabilized using EBioscience FOXP3 staining kit. Five x 10⁵ cells per tube were washed and incubated 45 minutes with anti-Hif1 α mAb at 5 μ g/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 0.13 log₁₀ fluorescent units when compared to a Mouse IgG1 negative control (Catalog #278-010). Binding was blocked when mAb was pre incubated with 100 μ g/ml recombinant Hif1 α .

* *Research Use Only. Not for use in Diagnostic procedures.*

Binding of anti-Hif1 α Mab +GAM/FITC to CoCl₂ treated fixed permeabilized Nalm-6 cells

