

PERFORMANCE DATA SHEET

2722

Monoclonal anti-human CD44*

mAb name/Clone: BU75

Isotype: Mouse IgG2a

Immunogen: Human tonsil germinal center B cell enriched

CATALOG#: 352-020

QUANTITY: 100 µg

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD44 molecules appear to serve as lymphocyte adhesion molecules with numerous functions in extracellular matrix binding, cell migration, hemopoiesis, and lymphocyte homing. Antibody BU75 recognizes epitope 2 of the CD44 molecule which interacts with hyaluronate (HA) mediating binding of lymphocytes to high endothelial venules. Antibody BU75 blocks binding of HA to CD44.

References: B.F. Hynes, et al, (1993) Adv Immunol **54**: 271-335. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford, (1995) p. 1713-1742. H.X. Liao, et al, (1995) J Immunol **155**: 3938-3945.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin.

PERFORMANCE: Five x 10⁵ cultured **Molt-4** human tumor cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD44 antibody at a concentration of 5 µg/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 1.75 log₁₀ fluorescent units when compared to a Mouse IgG2a negative control (Catalog #281-010).

Binding of anti-CD44(clone BU75) + GAM/FITC to human Molt-4 cells



* Research Use Only. Not for use in Diagnostic procedures.