

PERFORMANCE DATA SHEET

3143

Monoclonal anti-human CD134 (OX40)/R-PE *

mAb name/Clone: Ber Act 35

Isotype: Mouse IgG1

Immunogen: Human HUT 102 T cells

CATALOG#: 355-050

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2 ml

WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Human CD134 (OX40) is an activation-associated antigen which is predominantly expressed on activated CD4 positive cells. CD134 antigen is a member of the tumor necrosis factor (TNF) receptor family of molecules and may be involved with regulating T cell-dependent B cell proliferation and differentiation. Antibody Ber Act 35 recognizes the human CD134 molecule of about 35 kd.

References: U. Latza et al, Eur J Immunol (1994) **24**: 677-683. E. Stuber, et al, (1995) Immunity **2**: 507-521. Leukocyte Typing IV (W. Knapp, et al, eds.) Oxford University Press, Oxford, (1989) p. 464-465. Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995) p. 1157-1160.

STORAGE CONDITIONS: Store at 2 - 5°C. *Do not freeze! Protect from light.*

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at **0.25 mg/ml** with a A₅₆₅/A₂₈₀ ratio of 3.0.

PERFORMANCE: Five x 10⁵ cultured **HPB-MLT** human tumor cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD134/R-PE at a **1:50** dilution (5 µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.89 log₁₀** fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050). Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD134 antibody (Catalog #355-020).

**Research use only. Not for use in Diagnostic procedures.*

Binding of anti-CD134/PE to human HPB-MLT cells

