

PERFORMANCE DATA SHEET

3106

Monoclonal anti-human CD137 (4-1BB)/Biotin*

mAb name/Clone: 4B4-1

Isotype: Mouse IgG1 κ

Immunogen: Ectodomain of human 4-1BB fusion protein

CATALOG#: 360-030

QUANTITY: 100 μ g

CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD137 (4-1BB) is expressed on activated T cells within 24-48 hours of activation. CD137 is a type I membrane protein and a member of the tumor necrosis factor receptor superfamily. CD137 appears to be important for T cell proliferation and survival. Antibody 4B4-1 recognizes the CD137 molecule of about 35 kd from activated T cells.

References: M.R. Alderson, et al, (1994) Eur J Immunol **24**: 2219-2227. H.J. Gruss, and S.K. Dower, (1995) Blood **85**: 3378-3404. B.A. Garni-Wagner, et al, (1996) Cellular Immunol **169**: 91-98. H. Schwarz, et al, (1996) Blood **87**: 2839-2845

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/thawing not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Antibody from (low FBS containing) tissue culture supernatant was Protein A purified to >95% mouse immunoglobulin by SDS-PAGE (<1% bovine immunoglobulin), and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate using a desalting column.

PERFORMANCE: Cultured human CEM were stimulated 1 day in the presence of PMA (10ng/ml) and ionomycin (1 μ M). Five x 10⁵ cells were washed and incubated 45 minutes on ice with 80 μ l of anti-CD137/Biotin at 5 μ g/ml. Cells were washed twice and incubated with 2^o reagent Streptavidin/R-Phycoerythrin (Catalog #253-050), after which they were washed three times, fixed and analyzed by FACS. A 78% sub population of the cells stained positive with a mean shift of 1.95 log₁₀ fluorescent units when compared to a Mouse IgG1/Biotin negative control (Catalog #278-030). Binding was blocked when cells were pre incubated 10 minutes with 20 μ l of 0.5 mg/ml anti-CD137 antibody (Catalog #360-020).

*Research use only. Not for use in Diagnostic procedures.

Binding of anti-CD137/Biotin +SA/PE to stimulated human CEM cells

