

PERFORMANCE DATA SHEET

2439

Monoclonal anti-human CD357 (GITR AITR)/FITC*

mAb name/Clone: ANC5A3

Isotype: Mouse IgG3κ

Immunogen: Recombinant human GITR, PHA stimulated human PBL

CATALOG#: 368-040

QUANTITY: 120 tests

VOLUME IN VIAL: 0.2ml

WORKING DILUTION: (or use 1.6µl of concentrated stock per 5×10^5 -cell test)

INFORMATION: Glucocorticoid induced Tumor Necrosis Factor receptor family related gene (GITR, AITR) is a type 1 228 amino acid transmembrane protein that has been designated TNFRSF18. GITR is expressed mainly by CD4+ CD25+ T lymphocytes. Its engagement by antibodies or its ligand (GITRL) induces a costimulatory signal which can break T cell suppression (2). GITR has been implicated as a regulatory mechanism in anti-parasitic immune responses (4), and is involved with Macrophages activation in atherosclerotic plaque formation (3).

Clone ANC5A3 binds to human GITR in EIA and on stimulated cell surface in FACS. Preincubation of the receptor with an excess of this mAb will block binding of recombinant GITR Ligand.

References: 1) Gurney AL, et al. (1999) *Curr Biol* **9**(4): 215-218. 2) Ji HB, et al. (2004) *J Immunol* **172**(10): 5823-5827. 3) Kim WJ, WH Lee, et al. (2006) *Immunology* **119**(3): 44-9. 4) Furze RC, Selkirk ME, et al. (2006) *Microbes Infect* **8**(12-13): 2803-10. 5) Kohm AP, SD Miller, et al. (2004) *J Immunol* **172**: 4686-4690. 6) McHugh, R.S. et al. (2002) *Immunity* **16**:311. 7) Shin, H.H. et al. (2002) *FEBS Lett.* **514**:275. 8). Shimizu, J. et al. (2002) *Nature Immunol.* **3**:135.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein A purified antibody from tissue culture supernatant was reacted with FITC. Unconjugated FITC was separated from antibody/FITC conjugate by desalting column. The antibody/FITC conjugate is at **1.0 mg/ml** with a Fluorescein/IgG molar ratio of 5.2.

PERFORMANCE: Ficoll prepared human **peripheral blood mononuclear cells** were stimulated 5 days in culture with PHA. Five $\times 10^5$ cells per tube were washed and pre incubated 5 minutes with 20 µl of 250 µg/ml human IgG (to block non specific binding) after which they were incubated 45 minutes on ice with 80 µl of anti-GITR/FITC at a **1:50** dilution (20µg/ml). Cells were washed three times and analyzed by FACS. A net 24% sub population of the cells stained positive with a mean shift of **1.24 log₁₀** fluorescent units when compared to a Mouse IgG3/FITC negative control (Catalog #287-040). Binding was blocked when cells were pre incubated with a 10-fold excess of unlabeled anti-GITR mAb (cat #368-020).

***Research use only. Not for use in Diagnostic procedures.**

Binding of anti-GITR/FITC to stimulated human PBL

