## PERFORMANCE DATA SHEET

1818

## Monoclonal anti-human CD45RA\*



*mAb name/Clone:* 158.4D3 *Isotype:* Mouse IgG2a

Immunogen: Stimulated human leukocytes

**CATALOG#: 394-020 QUANTITY: 100 μg** 

CONCENTRATION: 1.0 mg/ml

**INFORMATION:** Human CD45RA is expressed on all cells of hematopoietic origin, except erythrocytes. CD45RA is a transmembrane tyrosine phosphate which can exist in at least nine different isoforms resulting from tissue-specific alternative RNA splicing of exons 4-7 of a single gene coding for the various N-terminal peptide segments. The CD45RA isoform predominates on naive/resting T cells and medullary thymocytes. Antibody 158.4D3 recognizes all CD45 molecules containing the A region exon (1,2).

*References:* 1) Leukocyte Typing III (A.J. McMichael, et al, eds.) Oxford University Press, Oxford, (1987) p. 788-817. 2) A. Gaya, et al, (1990) Intl Immunol 2:685-689. 3.) I.S. Trowbridge & M.L. Thomas, (1994) Annu Rev Immunol 12: 85-116.

**STORAGE CONDITIONS:** *Store at 2 - 5°C*. Freeze/Thawing is not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5 mg/ml Gentamicin Sulfate (as a preservative).

**PRODUCTION:** Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin.

**PERFORMANCE:** Five x  $10^5$  cultured human **Raji** cells were incubated 45 minutes on ice with 80  $\mu$ l of anti-CD45RA antibody at **5\mug/ml**. Cells were washed twice and incubated with  $2^0$  reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.86** log<sub>10</sub> fluorescent units when compared to a Mouse IgG2a negative control (Catalog #281-010).

\*Research use only. Not for use in Diagnostic procedures.

Binding of anti-CD45RA antibody to human cell lines

