

## PERFORMANCE DATA SHEET

1818

# Monoclonal anti-human CD45RA/R-PE\*

**mAb name/Clone:** 158.4D3

**Isotype:** Mouse IgG2aκ

**Immunogen:** Stimulated human leukocytes

**CATALOG#:** 394-050

**QUANTITY:** 120 tests

**VOLUME IN VIAL:** 0.2 ml

**WORKING DILUTION:** 1:50 (or use 1.6µl of concentrated stock per 5 x 10<sup>5</sup>-cell test)

**INFORMATION:** Human CD45RA is expressed on all cells of hematopoietic origin, except erythrocytes. CD45RA is a transmembrane tyrosine phosphate which can exist in at least nine different isoforms resulting from tissue-specific alternative RNA splicing of exons 4-7 of a single gene coding for the various N-terminal peptide segments. The CD45RA isoform predominates on naive/resting T cells and medullary thymocytes. Antibody 158.4D3 recognizes all CD45 molecules containing the A region exon (1,2).

**References:** 1) Leukocyte Typing III (A.J. McMichael, et al, eds.) Oxford University Press, Oxford, (1987) p. 788-817. 2) A. Gaya, et al, (1990) Intl Immunol 2:685-689. 3.) I.S. Trowbridge & M.L. Thomas, (1994) Annu Rev Immunol 12: 85-116.

**STORAGE CONDITIONS:** Store at 2 - 5°C. Do not freeze! Protect from light.


**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: \_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Protein A purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at 250 µg/ml with an A<sub>565</sub>/A<sub>280</sub> ratio of 3.28.

**PERFORMANCE:** Five x 10<sup>5</sup> cultured human **Raji** cells per tube were incubated 45 minutes on ice with 80 µl of anti-CD45RA/R-PE at a dilution factor of 1:50 (5µg/ml). Cells were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of 2.61 log<sub>10</sub> fluorescent units when compared to a Mouse IgG1/R-PE negative control (Catalog #278-050) at a similar concentration. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-CD45RA antibody (Catalog #394-020).

Binding of anti-CD45RA/R-PE to human Raji cells



**\*This Product is intended for Laboratory Research use only.**

R-Phycoerythrin (R-PE) is covered under patents: U.S. 4,520,110; European 76,695 and Canadian 1,179,942.