

## Human CD95(APO-1/FAS)-huIg *Fusion Protein\**

**CATALOG#:** 506-020

**QUANTITY:** 25 µg

**CONCENTRATION:** 0.5 mg/ml

**Molecular Structure:** A soluble fusion protein consisting of the mature extracellular (159 aa) domain of human CD95 fused to a linker (4 aa) and human IgG1 Fc + hinge (232 aa), with a predicted monomeric molecular weight of 44.3 kd.

**Transfectant Cell Line:** CHO

**INFORMATION:** Human CD95 (APO-1/FAS) is a type I cell surface glycoprotein that is strongly upregulated on activated T cells, B cells, NK cells and thymocytes (1). CD95 plays an important role in programmed cell death or apoptosis (2). Apoptosis appears to be a mechanism for regulating the immune response (3, 4). CD95-huIg fusion protein blocks binding of anti-human CD95 antibody as well as recombinant CD95L to cells expressing CD95. **References:** (1). Leukocyte Typing V (S.F. Schlossman, et al, eds.) Oxford University Press, Oxford (1995)

p. 1142-1148. (2). S. Nagata & P. Golstein (1995) Science 267: 1449-1456. (3). S. Nagata & T. Suda (1995) Immunol Today 16: 39-43. (4). D.H. Lynch, F. Ramsdell & M.R. Alderson (1995) Immunol Today 16: 569-574.

**STORAGE CONDITIONS:** Store at 2 - 5°C. Freeze/Thawing is not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: \_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 0.5mg/ml Gentamicin Sulfate (as a preservative).

**PRODUCTION:** CD95-huIg fusion protein was Protein A purified from (low FBS containing) tissue culture supernatant of CHO transfectants. Purity was >95% by SDS-PAGE with less than 1% Bovine Immunoglobulin.

**PERFORMANCE:** CD95-huIg fusion protein was reactive in EIA using a Goat-anti-human-Ig antibody capture, and detection by anti-CD95/Biotin (Cat # 316-030) and Streptavidin/HRP.

*\* Research Use Only. Not for use in Diagnostic procedures.*