## PERFORMANCE DATA SHEET

3223

## Human CD269(BCMA)-muIg/Biotin Fusion Protein\*



CATALOG#: 519-030 QUANTITY: 25 μg CONCENTRATION: 0.5 mg/ml

**Molecular Structure:** A soluble molecule consisting of the extracellular (54aa) domain of human BCMA fused to the

murine IgG2a Fc (232 aa). Predicted monomeric weight: 33 kd.

Transfectant Cell Line: CHO

INFORMATION: The human B cell activating factor (BAFF) and APRIL(a proliferation inducing ligand) are both type II molecules belonging to the TNF superfamily. They are expressed by non-B cells, and are down regulated by mitogenic stimulation(2). BAFF and APRIL bind to at least two receptors: TACI (transmembrane activator and CAML-interactor) and BCMA (B cell maturation antigen), both of which are restricted to B cells(3,4). Ligation of these receptors with recombinant BAFF dramatically increases IgM production by peripheral blood B cells(1). Recently a third receptor for BAFF (BAFF-R) was described(5). BAFF and BAFFR knockout mice have a reduced numbers of mature B cells in the periphery, however TACI and BCMA knockouts do not share this phenotype, suggesting that BAFF-R may the primary receptor for BAFF in mice(8,9,10). Cell surface BAFF can be proteolytically cleaved to form a soluble trimeric molecule(2). Levels of soluble BAFF correspond with levels of autoantibodies in Sjogren's Syndrome(11). Recombinant human BCMA-muIg binds to recombinant BAFF-muCD8 and can inhibit binding of this molecule to receptors on Raji cells.

References: 1) Schneider P., J. Tschopp, et al. *J. Exp. Med.* 1999, 189(11):1747-1756. 2) Shu, H.B., H. Johnson, W.H. Hui. *J Leukoc Biol* 1999, 65:680-683. 3) Marsters, S.A., A. Ashkenazi, et al. 2000, Curr Biol 10:785-788. 4) Xia, X., H. Hsu, et al. 2000, *J Exp Med.* 192(1): 137-143. 5) Thompson J.S., C. Ambrose, et al. Science 2001, 293: 2108-2111. 6)Roschke, V, T.S. Migone, et al. *J Immunol.* 2002, 169: 4314-4321. 7) MacLennan, C.M., C.G. Vinuesa, 2002, Immunity 17:235-238. 8) B. Schiemann, et al. (2001) Science 293: 2111-2114. 9) S.M. Harless, et al. (2001) Curr Biol 11: 1988-1989. 10) Mol Cell Biol (2001) 21: 4067-4074. 11) X. Mariette, et al. (2003) Ann Rhem Dis 62: 168-171.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended.


**PRODUCT STABILITY:** Product should retain activity for at least 6 months after shipping date when stored as recommended. Ship Date:

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl, 5% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Fusion protein from (low FBS containing) tissue culture supernatant of transfectants was purified using affinity and size exclusion chromatography and reacted with NHS-Biotin. Unconjugated Biotin was removed from conjugate by size exclusion chromatography.

**PERFORMANCE:** CD269(BCMA)-muIg/Biotin at 5 ug/ml bound to BAFF-coated CML beads in FACS using SA/PE detection. Beads stained positive with a **2.73** log shift compared to muIgFc Control Protein/Biotin (Cat # 581-030). Binding was blocked when beads were preincubated with 50-fold excess of unlabeled CD269(BCMA)-muIg (Cat # 519-020). An Irrelevant coated CML did not exhibit positive staining.

## Binding of CD269-mulg/Biotin +SA/PE to BAFF-muCD8 coated CML beads



<sup>\*</sup> Research Use Only. Not for use in Diagnostic procedures.