

Human CD127(IL-7R α)-muIg Fusion Protein*

CATALOG#: 539-820 (Preservate-Free)

QUANTITY: 25 μ g

CONCENTRATION: 0.5 mg/ml

Molecular Structure: A soluble molecule consisting of murine CD8 alpha signal peptide residual amino acids and linker:

(1)kpqapelrgs(10)

the mature extracellular domain of human CD127 (IL-7R α):(11)esgyaqngdledaelddysfscysqlevngsqhsltcafedpdvntnlfeicgalvevkcfnrklqeiyyfietkkflligksnicvkveksltckkidlttivkpeapf
dlsvvyregandfvvftntshlqkkyvkvlmhdvayrqekdenkwthvnlsstklflqrklqpaameyikvrsipdhfykgfwsewspysyfrtpeinnssgemd(29)

murine IgG2a Fc + hinge regions:

(230)gteprgptikpcppckcpapnllggpsvfifppkikdvlmislspivtcvvdvseddpdvqiswfvnnvevhtaqtqthredynstlrsvsalpiqhqdwmmsg
kefkckvnnkdlpapiertiskpkgsrapqvvyvlpppeeemtkkqvltcmvtdfmpediyvewtngktelnykntepvldsdgsyfmysklrvekknwvern
syscsvgheglhnhhtksfsrtpgk(464)

(464 aa total). The molecule is dimeric with a predicted monomeric non glycosylated molecular weight of 52.8 kd.

Transfectant Cell Line: CHO

INFORMATION: CD127 (IL-7R α) is the specific receptor component for the cytokine Interleukin -7 (IL-7). It is found on a wide variety of hematopoietic cell types including B cell precursors, and the majority of T cells. Its expression levels are decreased on T cells following activation. CD127 can dimerize with CD132(IL-2R γ) to form a high affinity IL-7 receptor (1). CD127 engagement is necessary for T cell development in humans (2).

References: 1) Noguchi, M. *et al.*, 1993, *Science* **262**:1877. 2) Pribyl, J.A. and T.W. LeBien, 1996, *Proc. Natl. Acad. Sci.* **93**:10348. 3) Goodwin RG, Friend D, Ziegler SF, et al. *Cell*. 1990;60:941.

STORAGE CONDITIONS: Store at 2 - 5°C. Open under aseptic conditions. Freeze/Thawing is not recommended.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date: _____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl. Product was sterile filtered under aseptic conditions.

PRODUCTION: Human CD127-muIg fusion protein was Protein A purified from (low FBS containing) tissue culture supernatant of CHO transfectants. Product was 0.2 μ sterile-filtered and vialled under aseptic conditions.

PERFORMANCE: Recombinant soluble CD127-muIg was reactive with anti-CD127 clone 40131 as well as clone ANC8F2 (catalog #348-030) in EIA and FACS.

**For Research Use Only. Not for Use in Diagnostic Procedures.*